



NCC Pediatrics Continuity Clinic Curriculum: **Sharing Unexpected News** *Faculty Version*

Goals & Objectives:

Goal: Increase your knowledge and skill in delivering difficult and unexpected news, and disclosing adverse events to families.

Objectives: At the end of this module, the pediatric resident should be able to:

- Recognize challenges specific to delivering unexpected news and disclosure of medical errors.
- Describe the elements of the mnemonics “SPIKES, NURSE, and GUIDE”
- Apply SPIKES, NURSE, or GUIDE to clinical scenarios that require delivering unexpected news or disclosure of medical errors.

Pre-Meeting Preparation:

- “Disclosure of Adverse Events in Pediatrics” (*AAP Policy Statement, Pediatrics, 2025*)
- "When Bad News Isn't Necessarily Bad: Recognizing Provider Bias When Sharing Unexpected News (*Carroll et al. Pediatrics, 2018*)
- Mnemonics and Frameworks (*VitalTalk, 2019*)
- Do's and Don'ts, SPIKES strategies for delivering unexpected news
- Watch the [Ted Talk](#) called "Doctors make mistakes. Can we talk about that?"
- Recall a situation where you had to share unexpected news and be prepared to discuss.

Conference Agenda:

- Review the sharing unexpected news quiz answers (*5 minutes*)
- Discuss your own case when you delivered unexpected news
- Practice delivering unexpected news using the scenarios included here. Make sure that you leave time for meaningful feedback. (*10 minute scenario and 5 minute feedback*)

Extra Credit:

- [Implementation Tools and Resources](#)
- [Medical Disclosure Powerpoint Slides from Ms. Barbara Moidel](#)
- ["SPIKES -- A Six-Step Protocol for Delivering Bad News: Application to the Patient with Cancer" \(The Oncologist, 2000\)](#)
- [“Disclosing Medical Mistakes: A Communication Management Plan for Physicians" \(Permanente Journal, 2013\)](#)
- [“Teaching Physicians How to Break Bad News” \(Archives of Pediatric and Adolescent Medicine, 1999\)](#)
- ["The Many Faces of Error Disclosure: A Common Set of Elements and a Definition" \(Society of General Internal Medicine, 2007\)](#)
- ["Medical Error Disclosure Among Pediatricians" \(Archives of Pediatric and Adolescent Medicine, 2008\)](#)
- [States with Apology Protection Laws](#)



Disclosure of Adverse Events in Pediatrics: Policy Statement

Laura Sigman, MD, JD, FAAP,¹ Robert Turbow, MD, JD, FAAP,² Daniel Neuspiel, MD, MPH, FAAP,³ Julia M. Kim, MD, MPH, FAAP,⁴ and the Committee on Medical Liability and Risk Management, and Council on Quality Improvement and Patient Safety

Disclosure of adverse events has become the expectation in medicine and is widely regarded as the appropriate path when medical errors occur. Although data are limited on adverse events in pediatrics, that they occur frequently is uncontested. Types and rates of errors vary depending on the care setting and patient population. Patients with complex medical conditions or from historically marginalized groups or minoritized communities likely suffer disparate health and safety outcomes. Systemic factors, including nonpunitive safety cultures and supportive environments within institutions, are essential to promoting disclosure. State laws protecting apologies from use in legal proceedings can also help to encourage open communication. Some states have adopted laws to advance disclosure, and governmental agencies provide materials encouraging open communication and early resolution after adverse events occur. Many programs emphasize the importance of supporting health care workers involved in adverse events. Shame, fear of professional and legal repercussions, and lack of training remain barriers to disclosure. Education for health care clinicians, support in health care settings, additional research on programs and disparities, and governmental and regulatory initiatives can support disclosure of adverse events.

INTRODUCTION

Over the last 2 decades, there has been increased recognition of the frequency and impact of adverse events in health care, accompanied by a transformation in how disclosure of such events takes place. Patients and their families expect such disclosure in a timely fashion. Additionally, data suggest that prompt and accurate disclosure of adverse events may be associated with decreased legal liability for the health care clinician. This policy statement updates a previous version¹; discusses the current

abstract

¹Armstrong Institute for Patient Safety and Quality, Department of Pediatrics, Johns Hopkins University School of Medicine, Baltimore, Maryland; ²Dignity Health- Central Coast California and Adjunct Professor Biomedical Engineering California Polytechnic State University, San Luis Obispo, California; ³Department of Pediatrics, Atrium Health; and ⁴Department of Pediatrics, Johns Hopkins University School of Medicine, Armstrong Institute for Patient Safety and Quality, Baltimore, Maryland

All authors contributed substantially to the conception and design; review and interpretation of relevant literature; drafting and revisions; and final approval of the version to the published.

CONFLICT OF INTEREST DISCLOSURE: The authors have indicated they have no potential conflicts of interest to disclose.

FUNDING: No external funding.

<https://doi.org/10.1542/peds.2025-070880>

Copyright © 2025 by the American Academy of Pediatrics

To cite: Sigman L, Turbow R, Neuspiel D, et al; American Academy of Pediatrics, Committee on Medical Liability and Risk Management, and Council on Quality Improvement and Patient Safety. Disclosure of Adverse Events in Pediatrics: Policy Statement. *Pediatrics*. 2025;155(4):e2025070880

| Table 1. Definitions | |
|-------------------------------------|--|
| Term | Definition |
| Patient safety | Prevention of patient harm and freedom from accidental injury in health care setting |
| Adverse event | Patient harm caused by medical care |
| Medical error | Act of commission or omission that unreasonably increases risk of an undesirable patient outcome |
| Preventable adverse event | Patient harm related to a medical error |
| Nonpreventable adverse event | Patient harm in absence of medical error |
| Potential adverse event (near miss) | Medical error with potential to cause patient harm that does not do so |
| Disclosure | Communication from health care personnel to affected patient and/or family about an adverse event |
| Reporting | Exchange of information about adverse events among clinicians and regulators |
| Just culture/safety culture | An organization's shared perceptions, beliefs, values, and attitudes that promote safety and minimize harm |

state of disclosure in pediatrics and issues surrounding it, including legal and ethical implications; and provides recommendations for pediatricians.

Patient safety is the prevention of patient harm and freedom from accidental injury in the health care setting.² An adverse event (AE) occurs when patient harm is caused by medical care.³ A medical error (ME) is an act of commission or omission that increases risk of an unintended patient outcome. AEs may be preventable (when patient harm is related to an ME) or nonpreventable (when patient harm occurs in the absence of an ME). An ME that causes patient harm becomes a preventable AE. An ME that has the potential to cause patient harm but does not do so is referred to as a potential AE or "near miss."⁴ These concepts and their relationships are explained in Table 1 and illustrated in Figure 1.

Estimates of total annual mortality in the United States attributable to preventable AEs vary widely, from 22 000⁵ to as high as 250 000 deaths.⁶ It is estimated that 6% of patients have experienced preventable harm.⁷ The American Academy of Pediatrics (AAP) has called attention to the importance of pediatric patient safety since 2001⁸ and has recommended improved identification and reporting of MEs and AEs to improve the culture of safety in pediatric care.⁹

The economic cost of AEs in the United States is substantial, although estimates have varied. In 2006, costs of AEs were estimated between \$393 and \$958 billion, equal to 18% to 45% of total US health care spending.¹⁰ In 2008, an estimate of the cost of MEs in the United States based on claims data was \$17.1 billion.¹¹ Another study estimated that MEs in the United States cost \$985 million in 2008 and over \$1 billion in 2009, with median costs per error of \$892 in 2008 and \$939 in 2009.¹² Specific costs for AEs involving children and adolescents have not been quantified.

ADVERSE EVENTS AND MEDICAL ERRORS IN PEDIATRIC POPULATIONS

While the exact magnitude of harm to pediatric patients from AEs is not well established, pediatric AEs occur frequently, with significant morbidity and mortality. Among Canadian inpatient pediatric AEs, the overall rate was 9.2%, and those related to surgery were most frequent¹³; almost half of these AEs were determined to be preventable. There are wide variations in estimates of the incidence of AEs among pediatric inpatients.^{9,14,15} Error rates related to medications in pediatrics have been noted to be 3 times higher than in adult patients.¹⁶

Types and rates of MEs vary depending on the health care setting and patient population. The majority of research to date has focused on errors in inpatient or emergency department settings. Emergency care settings involve high risks for errors, given the often busy, high-volume environment with frequent interruptions and transitions in care; complex patients who may not be well-known to staff; boarding; and commonly, a lack of standardized dosing, formulary, or information technology systems with pediatric safety features.¹⁷⁻¹⁹ Most pediatric care occurs in ambulatory settings, and studies have reported significant numbers of errors related to medications, vaccines, diagnoses, and coordination and transition of care.^{20,21} Medication administration errors account for the majority of preventable adverse drug events in the outpatient pediatric setting.²¹⁻²⁴ Although diagnostic errors occur frequently in




Figure 1.
Graphical Depiction of Adverse Events and Medical Errors

pediatrics, relatively little research has been performed on their incidence and epidemiology.^{25,26}

Children with special health care needs and/or medical complexity likely have higher rates of AEs than the general population because of frequent interactions with the health care system, clinicians in multiple settings, and medical needs. They are particularly vulnerable to medical errors during care transitions.^{24,27–32} Families are an underused resource for detecting and reporting safety events in these populations and can provide valuable perspectives, including in ambulatory settings.^{33–36}

Although research is limited on rates of AEs in historically marginalized or minoritized populations, evidence shows that these communities are more likely to suffer disparate health and safety outcomes.^{37,38} Recent data have shown higher rates of AEs in hospitalized Latino children and publicly insured children, including serious AEs.^{39,40} Hospitalized children of parents with limited comfort with English were twice as likely to experience AEs from medical care.⁴¹ Language barriers have been associated with increased rates of AEs^{42,43}; underreporting of AEs in hospitals relying on voluntary event reports⁴⁴; and less willingness to question health care clinicians, which may contribute to increased safety events.⁴⁵

THE SYSTEMS APPROACH AND SAFETY CULTURE

The Institute of Medicine (IOM) noted that most MEs are attributable to flaws in systems rather than individuals

and called for a “dramatic improvement in the reliability and safety” of the health care process.³ For this improvement to occur, AEs must first be identified and analyzed to understand their preventable causes and to allow for systematic safety improvements. Disclosure and open communication with patients and their families after an AE may benefit the patient and health care clinicians, reduce consequential harms, allow for better follow-up, and promote a safety culture.

Safety culture refers to an environment that facilitates open and honest communication to promote safety and minimize harm⁴⁶ and is particularly important when potentially uncomfortable topics such as MEs or preventable harm are being addressed. A growing body of literature, including subspecialty publications in pediatrics, demonstrate that safer culture is associated with safer care. Increasingly, the culture of pediatrics, and health care in general, is to promptly disclose to patients and families after an AE has taken place. Multiple factors are considered when disclosing MEs, including safety culture, clinician and patient level factors, and the type of error itself (see Figure 2).

REPORTING VS DISCLOSURE

The concepts of *reporting* and *disclosure* of AEs should be distinguished. *Reporting* refers to the exchange of information among clinicians and regulators. Reporting systems may be internal to health care organizations or may be




Figure 2. Influences on the Decision to Disclose a Medical Error (adapted from Fein et al⁹¹ [<https://www.ahrq.gov/downloads/pub/advances/vol2/Fein.pdf>])

required by licensing boards and governmental regulations. These systems may be voluntary or mandatory, depending on state and institutional policies, and some organizations may use automated AE reporting. Prior reporting systems often focused on punitive consequences, which was found to deter further reporting. More recent trends emphasize the adoption of “just culture,” promote reporting of AEs by all clinicians and staff, and utilize reports to develop systematic improvements and a safer patient care environment.^{47,48} Additionally, there is emerging evidence that including family reports in safety surveillance systems, which is not typical in many health care institutions, may help to identify more AEs.³⁴

DISCLOSURE TO PATIENTS AND FAMILIES

Historically, physicians were not often advised to disclose MEs to patients and their families, but early disclosure and resolution is now routinely encouraged. A variety of programs have been initiated around the United States, including the CANDOR (communication and optimal resolution) approach (<https://www.ahrq.gov/patient-safety/settings/hospital/candor/index.html>).⁴⁹ The CANDOR program consists of 2 main components:

1. Prompt and accurate disclosure to families
2. Care for the caregiver (see “Effects on Health Care Clinicians” section)

Clinicians involved are expected to notify their malpractice attorney(s) or institutional legal and risk management contacts as soon as possible. It is important for trainees to immediately notify appropriate supervisors and involve them, whenever possible, in conversations with families. Additional personnel, such as office managers, unit directors, and social workers, can aid in responding and provide support and guidance. Prompt and accurate discussion is encouraged, and whenever possible, a care team should have an organized plan, including predisclosure huddles and checklists, such as those available through the AHRQ, prior to meeting with families. In the immediate aftermath of a serious event, and when there may not be time to involve additional resources, conversations with families are usually limited to statements that something went wrong and that a formal investigation—usually in the form of a “root cause analysis” or “apparent cause analysis”—is forthcoming. At the initial discussion, clinicians can advise families that not all the facts are known and can assure them that additional meetings will be scheduled as more information is available. Apologies, including statements of sympathy with or without expressions of fault, may be included in initial conversations depending on the institutions and clinicians involved as well as the legal landscape around disclosures and protections for apologies and statements of fault in the local jurisdiction.^{50,51} A more formal

disclosure is usually done after an investigation has been conducted, and at times in conjunction with an early offer for resolution. Attorneys or risk management staff usually direct such discussions. Veracity, meaning that communication should be honest, is one of the guiding principles of bioethics and a moral foundation of disclosure. The principle of truth is deeply rooted in the practice of medicine and is essential for building trust between clinicians and patients or families. Although in the past, patients and their families were often “shielded” from the truth by paternalistic physicians, they should be viewed as partners in shared decision making. Families cannot make informed decisions for their loved ones if they have not been given relevant facts.

ENGAGEMENT OF CHILDREN AND ADOLESCENTS

Children and adolescents, including those with chronic illnesses, have expressed their desire to be involved in disclosures of errors affecting their care,⁵² and consideration should be given to involving them in such discussions when appropriate. Disclosure to children should be individualized and should not be determined simply by age and developmental parameters. Koller et al⁵³ have suggested that the development of policies on disclosure “must begin by examining children’s understanding of medical errors and what they expect from their health care clinicians when errors occur.”

ADOLESCENT CONFIDENTIALITY

There may be circumstances when AEs occur during the confidential treatment of adolescents, including conditions involving behavioral health, substance use, or sexual activity. Although state laws vary depending on exact age and condition, adolescents are usually legally entitled to seek treatment without needing consent from a parent or guardian for these conditions.⁵⁴ In such cases, disclosures are limited to the adolescent alone. Clinicians need to remain cognizant that subsequent care for medical consequences of the AE may fall outside adolescent confidentiality protections, and state laws protecting parental rights may impose further complications. Consultation with legal and/or risk management advisors in the relevant jurisdiction is recommended because of the complexity and variety of these protections.

BEST PRACTICES FOR DISCLOSURE

Disclosure practices may vary depending on setting, type of error, and patient population. Important aspects of disclosure include an explanation of what happened, acknowledgment that something has gone wrong and of responsibility, expression of sincere regret and apology, and commitment to preventing recurrences.⁵⁵ These should be conducted with the involvement of appropriate parties in each practice

setting. Participants may include leaders, risk managers, and possibly attorneys. Immediately after an AE occurs, clinicians can follow recommended steps such as involving team members, practice or unit leaders, risk managers, or attorneys; apologizing when appropriate; and communicating with the involved patient and/or family. Resources for responding to AEs are provided in the Toolkit Appendix.

EFFECTS ON HEALTH CARE CLINICIANS

AEs and MEs not only affect patients and their families but also may have devastating effects on health care clinicians, who may suffer emotional consequences both from preventable AEs and from subsequent malpractice litigation.^{56–58} Affected clinicians may feel guilt, shame, and isolation, and these feelings may be exacerbated by negative reactions from their colleagues. They may experience depression, anxiety, and/or posttraumatic stress disorder.⁵⁹ Anticipated or actual punitive consequences can add further emotional and financial burdens on clinicians. Although the terminology is evolving, the concepts of “second victims” or more recently “care for the caregiver” acknowledge that clinicians often experience significant personal turmoil if an AE occurs and focus on the important need to support health care clinicians involved in these events. Support systems for affected clinicians may reduce distress among health care clinicians.^{60–62} The tragedy in 2011 of a pediatric intensive care unit nurse’s suicide following a medication error highlights the devastating effects that MEs can have on the health care workers as well.⁵⁹

BARRIERS TO AND PROMOTERS OF DISCLOSURE

Numerous barriers to disclosure have been identified, including shame, fear of litigation and punishment, concern about the impact on professional reputation, decreased patient trust, situational complexity, lack of training or confidence in how to disclose, inconsistent guidelines on disclosure, and the lack of a nonpunitive patient safety culture.⁶³ Language and cultural barriers may hinder communication about AEs. Even with institutional policies advocating for disclosure, errors may not always be disclosed. In some cases, failure to disclose may impact patients at other institutions. For example, an outbreak of carbapenem-resistant *Enterobacter* infections related to duodenoscope contamination at one institution was associated with later deaths in other hospitals and states.⁶⁴

Factors that promote disclosure include a safe setting for reporting AEs and availability of guidelines and education on how to disclose errors.^{65,66} Peer support for disclosure, just-in-time disclosure coaching, refresher trainings, medical school and residency disclosure training and modeling, and organizational and national policies supporting disclosure also influence the likelihood and implementation of disclosure.⁵⁵

LEGAL RISKS AND OUTCOMES

Patients and caregivers most often desire complete disclosure of AEs.^{67,68} Yet historically, lawyers advised their physician-clients not to disclose MEs and did so with sound legal justification, because of the risk of statements of apology being used against physicians if an AE should result in a lawsuit.^{51,69,70} Physicians were trained that admitting fault would increase the risk of being sued.

However, studies suggest that affected patients and families may be less likely to pursue litigation against their health care clinicians if such disclosure is provided.^{71–76} Lack of such communication may make patients feel worse and may erode the sense of trust in their caregivers that is key to healing and to optimal health care. Although there is some disagreement in the legal literature on the impact of apologies and admissions of fault,^{77–79} the preponderance of the data, as well as the guiding ethical principles, support prompt and accurate disclosure.⁸⁰ Patients and families expect information about the error, sincere remorse, and a pledge of improvement.⁶⁶

Numerous states have passed so-called “apology laws,” which protect statements of apologies from being used in court against health care clinicians. As of 2022, 39 states and the District of Columbia have such statutes.⁸¹ Most of these states protect only sympathetic statements, such as saying “I’m sorry for your suffering” or “I regret that this happened,” while 9 states also protect statements of fault.⁵⁰ These apology laws, along with institutional, state, and federal efforts to implement disclosure programs, may help to encourage candid discussions after AEs.⁷⁴ More recently, several states have developed “candor laws” that provide a process for investigating and communicating openly about AEs. Under the Colorado Candor Act, for example, discussions and offers of compensation held in the process are privileged and confidential, meaning they cannot be used against clinicians if the injured party later decides to file a lawsuit.⁸²

A case in which a nurse was convicted of negligent homicide after a medication error highlights the need for broader legal protections for statements about AEs. In this case in Tennessee, which was decided in 2022, the nurse’s words to state investigators were used against her in court. Widespread concern has been expressed that health care workers will be hesitant to disclose errors if their cooperation will be used as part of criminal prosecutions.^{83,84} Although this type of legal case is extremely rare, the potential deterring effect on self-reporting could not only undermine the disclosure of AEs but also reverse gains in establishing a culture of safety and other patient safety initiatives.⁸⁵

EDUCATION ON DISCLOSURE OF PREVENTABLE ADVERSE EVENTS

In 2019, disclosure of AEs was included as a common program requirement for resident education and experience,

highlighting the importance of disclosure as a patient safety skill by the Accreditation Council for Graduate Medical Education (ACGME) (<https://www.acgme.org/What-We-Do/Accreditation/Common-Program-Requirements>). With increased focus on experiential learning, trainees have been involved in real or simulation scenarios, including disclosure of AEs.^{86,87} Disclosure practices are also included as one of 13 core professional activities for medical school graduates by the Association of American Medical Colleges, as part of identifying system failures and contributing to a culture of safety and improvement.^{88,89} Continued education in patient safety and developing communication skills for effective disclosure is needed at all training and career levels.^{55,90}

CONCLUSION

Progress has been made toward routine disclosure of adverse events over the past decades. Further efforts to better understand adverse events in pediatrics, focusing on risks, different health care settings, and inequitable practices that disproportionately impact historically marginalized or minoritized populations, may help to improve trends in disclosure. Support from institutions and legal protections can facilitate a culture of safety, reduce fear of disclosure, and provide enhanced education and support for health care clinicians involved in adverse events.

RECOMMENDATIONS

For Pediatric Health Care Clinicians, Practices, and Institutions

1. Develop and implement policies and procedures for identifying and disclosing AEs to patients and families in an honest and empathetic manner as part of a nonpunitive safety culture.
2. Develop policies and procedures and provide resources to support clinicians and other staff involved in AEs.
3. Encourage a culture of safety, just culture, and reporting by all staff as well as by patients and families.
4. Identify populations and situations with higher risk for AEs, such as patients with chronic illnesses and those from historically marginalized or minoritized communities, and partner with families and care teams to help prevent them.

For Medical Educators

5. Develop and implement educational programs regarding identification and prevention of MEs and communication about AEs with patients and their families as part of a comprehensive patient safety curriculum.

For Researchers

6. Investigate the consequences of various approaches to disclosure as well as of the effectiveness of disclosure education.
7. Further explore and address disparities in AEs and disclosure. More data are needed on trends in AEs and disclosure by race, ethnicity, chronic conditions, and preferred language; impact of AEs and disclosure on historically marginalized or minoritized communities; structural factors related to disclosure; and impact on clinicians underrepresented in medicine.

For Pediatric Advocates

8. Encourage states and the federal government to adopt laws protecting apologies, programs supporting disclosure, and other mechanisms to reduce liability risks associated with disclosure in order to avoid detrimental effects on health care clinicians' reactions to AEs.

For more information about advocating for these issues, contact the AAP state advocacy team at stgov@aap.org.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

LEAD AUTHORS

Laura Sigman, MD, JD, FAAP
Robert Turbow, MD, JD, FAAP
Daniel Neuspiel, MD, MPH, FAAP
Julia M. Kim, MD, MPH, FAAP

COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT, 2023-2024

Steven A. Bondi, JD, MD, FAAP, Chairperson
Craig Howard Gosdin, MD, FAAP

David A. Horowitz, MD, FAAP
 Kajal Khanna, MD, JD, FAAP
 Jonathan Muraskas, MD, FAAP
 Alan L. Nager, MD, MHA, FAAP
 Sarah W. Northrop, MD, FAAP
 Preethi Srinivasakumar, MD, FAAP
 Michelle Terry, MD, FAAP
 Ryan Earl Alanzalon, MD, FAAP

COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT, PAST CONTRIBUTING MEMBERS

James Peter Scibilia, MD, FAAP, Immediate Past Chairperson
 Laura Sigman, MD, JD, FAAP
 Robert Turbow, MD, JD, FAAP

STAFF

Julie Kersten Ake
 Sunnah Kim

COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY, 2023-2024

Ulfat Shaikh, MD, FAAP Chairperson
 Francisco Javier Alvarez, MD, FAAP
 John Chuo, MD, FAAP
 Meghan Drayton Jackson, DO, FAAP
 Julia M. Kim, MD, FAAP
 Mary Beth Miotto, MD, FAAP
 Raina Paul, MD, FAAP
 Corinna Rea, MD, FAAP
 Sandra Spencer, MD, FAAP
 Amy Tyler, MD, FAAP
 Joyee Vachani, MD, FAAP

STAFF

Cathleen Guch

ABBREVIATIONS

AAP: American Academy of Pediatrics
 AEs: adverse events
 AHRQ: Agency for Healthcare Research and Quality
 IOM: Institute of Medicine
 MEs: medical errors.

REFERENCES



- McDonnell WM, Altman RL, Bondi SA, et al; COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT; COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY. Disclosure of adverse events in pediatrics. *Pediatrics*. 2016;138(6):e20163215. PubMed doi: 10.1542/peds.2016-3215

- Institute of Medicine, Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. National Academy Press; 2001.
- Institute of Medicine, Committee on Quality of Health Care in America. *To Err Is Human: Building a Safer Health System*. Kohn LT, Corrigan JM, Donaldson MS, eds. National Academies Press; 2000.
- Rodziewicz TL, Houseman B, Hipskin J. *Medical Error Reduction and Prevention*. StatPearls Publishing; 2022.
- Rodwin BA, Bilan VP, Merchant NB, et al. Rate of preventable mortality in hospitalized patients: a systematic review and meta-analysis. *J Gen Intern Med*. 2020;35(7):2099–2106. PubMed doi: 10.1007/s11606-019-05592-5
- Makary MA, Daniel M. Medical error-the third leading cause of death in the US. *BMJ*. 2016;353:i2139. PubMed doi: 10.1136/bmj.i2139
- Panagioti M, Khan K, Keers RN, et al. Prevalence, severity, and nature of preventable patient harm across medical care settings: systematic review and meta-analysis. *BMJ*. 2019;366:l4185. PubMed doi: 10.1136/bmj.l4185
- Lannon CM, Coven BJ, Lane France F, et al; National Initiative for Children's Health Care Quality Project Advisory Committee. Principles of patient safety in pediatrics. *Pediatrics*. 2001;107(6):1473–1475. PubMed doi: 10.1542/peds.107.6.1473
- Mueller BU, Neuspiel DR, Fisher ERS, et al; COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY, COMMITTEE ON HOSPITAL CARE. Policy Statement. Principles of pediatric patient safety: reducing harm due to medical care. *Pediatrics*. 2019;143(2):e20183649. PubMed doi: 10.1542/peds.2018-3649
- Goodman JC, Villarreal P, Jones B. The social cost of adverse medical events, and what we can do about it. *Health Aff (Millwood)*. 2011;30(4):590–595. PubMed doi: 10.1377/hlthaff.2010.1256
- Van Den Bos J, Rustagi K, Gray T, Halford M, Ziemkiewicz E, Shreve J. The \$17.1 billion problem: the annual cost of measurable medical errors. *Health Aff (Millwood)*. 2011;30(4):596–603. PubMed doi: 10.1377/hlthaff.2011.0084
- David G, Gunnarsson CL, Waters HC, Horblyuk R, Kaplan HS. Economic measurement of medical errors using a hospital claims database. *Value Health*. 2013;16(2):305–310. PubMed doi: 10.1016/j.jval.2012.11.010
- Matlow AG, Baker GR, Flintoft V, et al. Adverse events among children in Canadian hospitals: the Canadian Paediatric Adverse Events Study. *CMAJ*. 2012;184(13):E709–E718. PubMed doi: 10.1503/cmaj.112153
- Gates PJ, Meyerson SA, Baysari MT, Lehmann CU, Westbrook JL. Preventable adverse drug events among inpatients: a systematic review. *Pediatrics*. 2018;142(3):e20180805. PubMed doi: 10.1542/peds.2018-0805
- Schroeder L, Stockwell DC. The persistent challenge of understanding preventable adverse drug events. *Pediatrics*. 2018;142(3):e20181841. PubMed doi: 10.1542/peds.2018-1841

16. Kaushal R, Bates DW, Landrigan C, et al. Medication errors and adverse drug events in pediatric inpatients. *JAMA*. 2001;285(16):2114–2120. PubMed doi: 10.1001/jama.285.16.2114
17. Benjamin L, Frush K, Shaw K, et al; AMERICAN ACADEMY OF PEDIATRICS Committee on Pediatric Emergency Medicine; AMERICAN COLLEGE OF EMERGENCY PHYSICIANS Pediatric Emergency Medicine Committee; EMERGENCY NURSES ASSOCIATION Pediatric Emergency Medicine Committee. Pediatric medication safety in the emergency department. *Pediatrics*. 2018;141(3):e20174066. PubMed doi: 10.1542/peds.2017-4066
18. Gross TK, Lane NE, Timm NL, et al; COMMITTEE ON PEDIATRIC EMERGENCY MEDICINE. Crowding in the emergency department: challenges and best practices for the care of children. *Pediatrics*. 2023;151(3):e2022060972. PubMed doi: 10.1542/peds.2022-060972
19. Joseph MM, Mahajan P, Snow SK, Ku BC, Saidinejad M; AMERICAN ACADEMY OF PEDIATRICS COMMITTEE ON PEDIATRIC EMERGENCY MEDICINE, AMERICAN COLLEGE OF EMERGENCY PHYSICIANS PEDIATRIC EMERGENCY MEDICINE COMMITTEE, and EMERGENCY NURSES ASSOCIATION PEDIATRIC COMMITTEE. Optimizing pediatric patient safety in the emergency care setting. *Pediatrics*. 2022;150(5):e2022059673. PubMed doi: 10.1542/peds.2022-059673
20. Neuspiel DR, Stubbs EH. Patient safety in ambulatory care. *Pediatr Clin North Am*. 2012;59(6):1341–1354. PubMed doi: 10.1016/j.pcl.2012.08.006
21. Huth K, Hotz A, Starmer AJ. Patient safety in ambulatory pediatrics. *Curr Treat Options Pediatr*. 2020;6(4):350–365. PubMed doi: 10.1007/s40746-020-00213-4
22. Kaushal R, Goldmann DA, Keohane CA, et al. Adverse drug events in pediatric outpatients. *Ambul Pediatr*. 2007;7(5):383–389. PubMed doi: 10.1016/j.ambp.2007.05.005
23. Yin HS, Parker RM, Sanders LM, et al. Liquid medication errors and dosing tools: a randomized controlled experiment. *Pediatrics*. 2016;138(4):e20160357. PubMed doi: 10.1542/peds.2016-0357
24. Yin HS, Neuspiel DR, Paul IM, et al; American Academy of Pediatrics, Council on Quality Improvement and Patient Safety, Committee on Drugs. Policy statement. Preventing home medication administration errors. *Pediatrics*. 2021;148(6):e2021054666. PubMed doi: 10.1542/peds.2021-054666
25. Rinke ML, Singh H, Heo M, et al. Diagnostic errors in primary care pediatrics: Project RedDE. *Acad Pediatr*. 2018;18(2):220–227. PubMed doi: 10.1016/j.acap.2017.08.005
26. Marshall TL, Rinke ML, Olson APJ, Brady PW. Diagnostic error in pediatrics: a narrative review. *Pediatrics*. 2022;149(suppl 3):e2020045948D. PubMed doi: 10.1542/peds.2020-045948D
27. Rinke ML, Bundy DG, Shore AD, Colantuoni E, Morlock LL, Miller MR. Pediatric antidepressant medication errors in a national error reporting database. *J Dev Behav Pediatr*. 2010;31(2):129–136. PubMed doi: 10.1097/DBP.0b013e3181ce6509
28. Taylor JA, Winter L, Geyer LJ, Hawkins DS. Oral outpatient chemotherapy medication errors in children with acute lymphoblastic leukemia. *Cancer*. 2006;107(6):1400–1406. PubMed doi: 10.1002/cncr.22131
29. Walsh KE, Mazor KM, Stille CJ, et al. Medication errors in the homes of children with chronic conditions. *Arch Dis Child*. 2011;96(6):581–586. PubMed doi: 10.1136/adc.2010.204479
30. Stone BL, Boehme S, Mundorff MB, Maloney CG, Srivastava R. Hospital admission medication reconciliation in medically complex children: an observational study. *Arch Dis Child*. 2010;95(4):250–255. PubMed doi: 10.1136/adc.2009.167528
31. Feinstein JA, Feudtner C, Kempe A. Adverse drug event-related emergency department visits associated with complex chronic conditions. *Pediatrics*. 2014;133(6):e1575–e1585. PubMed doi: 10.1542/peds.2013-3060
32. Huth K, Vandecruys P, Orkin J, Patel H. Medication Safety for children with medical complexity. Canadian Pediatric Society. *Paediatr Child Health*. 2020;25(7):473–474. doi: 10.1093/pch/pxaa105
33. Khan A, Baird J, Kelly MM, et al. Family safety reporting in medically complex children: parent, staff, and leader perspectives. *Pediatrics*. 2022;149(6):e2021053913. PubMed doi: 10.1542/peds.2021-053913
34. Khan A, Furtak SL, Melvin P, Rogers JE, Schuster MA, Landrigan CP. Parent-reported errors and adverse events in hospitalized children. *JAMA Pediatr*. 2016;170(4):e154608. PubMed doi: 10.1001/jamapediatrics.2015.4608
35. Daniels JP, Hunc K, Cochrane DD, et al. Identification by families of pediatric adverse events and near misses overlooked by health care providers. *CMAJ*. 2012;184(1):29–34. PubMed doi: 10.1503/cmaj.110393
36. Lang S, Garrido MV, Heintze C. Patients' views of adverse events in primary and ambulatory care: a systematic review to assess methods and the content of what patients consider to be adverse events. *BMC Fam Pract*. 2016;17:6. PubMed doi: 10.1186/s12875-016-0408-0
37. Jean-Pierre P. Medical error and vulnerable communities. *Boston Univ Law Rev*. 2022;102:327–392.
38. Raman J, Johnson TJ, Hayes K, Balamuth F. Racial differences in sepsis recognition in the emergency department. *Pediatrics*. 2019;144(4):e20190348. PubMed doi: 10.1542/peds.2019-0348
39. Stockwell DC, Landrigan CP, Toomey SL, et al; GAPPs Study Group. Racial, ethnic, and socioeconomic disparities in patient safety events for hospitalized children. *Hosp Pediatr*. 2019;9(1):1–5. PubMed doi: 10.1542/hpeds.2018-0131
40. Lion KC, Rafton SA, Shafii J, et al. Association between language, serious adverse events, and length of stay among hospitalized children. *Hosp Pediatr*. 2013;3(3):219–225. PubMed doi: 10.1542/hpeds.2012-0091
41. Khan A, Yin HS, Brach C, et al; Patient and Family Centered I-PASS Health Literacy Subcommittee. Patient and Family Centered I-PASS Health Literacy Subcommittee. Association between parent comfort with English and adverse events among hospitalized children. *JAMA Pediatr*. 2020;174(12):e203215. PubMed doi: 10.1001/jamapediatrics.2020.3215
42. Cohen AL, Rivara F, Marcuse EK, McPhillips H, Davis R. Are language barriers associated with serious medical events in hospitalized pediatric patients? *Pediatrics*. 2005;116(3):575–579. PubMed doi: 10.1542/peds.2005-0521

43. Divi C, Koss RG, Schmaltz SP, Loeb JM. Language proficiency and adverse events in US hospitals: a pilot study. *Int J Qual Health Care*. 2007;19(2):60–67. PubMed doi: 10.1093/intqhc/mzl069
44. Halvorson EE, Thurtle DP, Easter A, Lovato J, Stockwell D. Disparities in adverse event reporting for hospitalized children. *J Patient Saf*. 2022;18(6):e928–e933. PubMed doi: 10.1097/PTS.0000000000001049
45. Khan A, Parente V, Baird JD, et al; Patient and Family Centered I-PASS SCORE Scientific Oversight Committee. Association of patient and family reports of hospital safety climate with language proficiency in the US. *JAMA Pediatr*. 2022;176(8):776–786. PubMed doi: 10.1001/jamapediatrics.2022.1831
46. Weaver SJ, Lubomski LH, et al. Promoting a culture of safety as a patient safety strategy: a systematic review. *Ann Int Med*. 2013;158(5.2):369–374. PubMed doi: 10.7326/0003-4819-158-5-2013051-00002
47. Boysen PG II. Just culture: a foundation for balanced accountability and patient safety. *Ochsner J*. 2013;13(3):400–406. PubMed
48. Marx D. Patient safety and the just culture. *Obstet Gynecol Clin North Am*. 2019;46(2):239–245. PubMed doi: 10.1016/j.ogc.2019.01.003
49. Agency for Healthcare Research and Quality. Communication and Optimal Resolution (CANDOR). Accessed July 19, 2024. <https://www.ahrq.gov/patient-safety/settings/hospital/candor/index.html>
50. Ross NE, Newman WJ. The role of apology laws in medical malpractice. *J Am Acad Psychiatry Law*. 2021;49(3):406–414. PubMed
51. Mastroianni AC, Mello MM, Sommer S, Hardy M, Gallagher TH. The flaws in state ‘apology’ and ‘disclosure’ laws dilute their intended impact on malpractice suits. *Health Aff (Millwood)*. 2010;29(9):1611–1619. PubMed doi: 10.1377/hlthaff.2009.0134
52. Koller D, Binder MJ, Alexander S, Darch J. “Everybody makes mistakes”: children’s views on medical errors and disclosure. *J Pediatr Nurs*. 2019;49:1–9. PubMed doi: 10.1016/j.pedn.2019.07.014
53. Koller D, Rummens A, Le Pouesard M, et al. Patient disclosure of medical errors in paediatrics: A systematic literature review. *Paediatr Child Health*. 2016;21(4):e32–e38. PubMed doi: 10.1093/pch/21.4.e32
54. Chung RJ, Lee JB, Hackell JM, et al; COMMITTEE ON ADOLESCENCE; COMMITTEE ON PRACTICE & AMBULATORY MEDICINE. Confidentiality in the care of adolescents: policy statement. *Pediatrics*. 2024;153(5):e2024066326. PubMed doi: 10.1542/peds.2024-066326
55. Wu AW, Boyle DJ, Wallace G, Mazor KM. Disclosure of adverse events in the United States and Canada: an update, and a proposed framework for improvement. *J Public Health Res*. 2013;2(3):e32. PubMed doi: 10.4081/jphr.2013.e32
56. Wu AW. Medical error: the second victim. The doctor who makes the mistake needs help too. *BMJ*. 2000;320(7237):726–727. PubMed doi: 10.1136/bmj.320.7237.726
57. Seys D, Wu AW, Van Gerven E, et al. Health care professionals as second victims after adverse events: a systematic review. *Eval Health Prof*. 2013;36(2):135–162. PubMed doi: 10.1177/0163278712458918
58. Coughlan B, Powell D, Higgins MF. The second victim: a review. *Eur J Obstet Gynecol Reprod Biol*. 2017;213:11–16. PubMed doi: 10.1016/j.ejogrb.2017.04.002
59. Grissinger M. Too many abandon the “second victims” of medical errors. *P&T*. 2014;39(9):591–592. PubMed
60. Edrees H, Connors C, Paine L, Norvell M, Taylor H, Wu AW. Implementing the RISE second victim support programme at the Johns Hopkins Hospital: a case study. *BMJ Open*. 2016;6(9):e011708. PubMed doi: 10.1136/bmjopen-2016-011708
61. Busch IM, Moretti F, Campagna I, et al. Promoting the psychological well-being of healthcare providers facing the burden of adverse events: a systematic review of second victim support resources. *Int J Environ Res Public Health*. 2021;18(10):5080. PubMed doi: 10.3390/ijerph18105080
62. Shapiro J, Galowitz P. Peer support for clinicians: a programmatic approach. *Acad Med*. 2016;91(9):1200–1204. PubMed doi: 10.1097/ACM.0000000000001297
63. Edwards MT. An assessment of the impact of just culture on quality and safety in US hospitals. *Am J Med Qual*. 2018;33(5):502–508. PubMed doi: 10.1177/1062860618768057
64. Centers for Disease Control and Prevention. Los Angeles County/UCLA investigation of CRE transmission and duodenoscopes. July 10, 2015. Accessed July 19, 2024. https://archive.cdc.gov/www_cdc.gov/hai/outbreaks/cdcstatement-la-cre.html#:~:text=In%20this%20investigation%2C%20exposure%20to,associated%20with%20transmission%20of%20CRE
65. Ock M, Lim SY, Jo MW, Lee SI. Frequency, expected effects, obstacles, and facilitators of disclosure of patient safety incidents: a systematic review. *J Prev Med Public Health*. 2017;50(2):68–82. PubMed doi: 10.3961/jpmph.16.105
66. Sattar R, Johnson J, Lawton R. The views and experiences of patients and health-care professionals on the disclosure of adverse events: A systematic review and qualitative meta-ethnographic synthesis. *Health Expect*. 2020;23(3):571–583. PubMed doi: 10.1111/hex.13029
67. Coffey M, Espin S, Hahmann T, et al. Parent preference for medical error disclosure: a qualitative study. *Hosp Pediatr*. 2017;7(1):24–30. PubMed doi: 10.1542/hpeds.2016-0048
68. Hobgood C, Tamayo-Sarver JH, Elms A, Weiner B. Parental preferences for error disclosure, reporting, and legal action after medical error in the care of their children. *Pediatrics*. 2005;116(6):1276–1286. PubMed doi: 10.1542/peds.2005-0946
69. Hyman DA. When and why lawyers are the problem. *DePaul L Rev*. 2008;57(2):267–280.
70. Tabler N. Should physicians apologize for medical errors? *Health Lawyer*. 2007;19:23–30.
71. Helmchen LA, Richards MR, McDonald TB. How does routine disclosure of medical error affect patients’ propensity to sue and their assessment of provider quality? Evidence from survey data. *Med Care*. 2010;48(11):955–961. PubMed doi: 10.1097/MLR.0b013e3181eaf84d
72. Kachalia A, Kaufman SR, Boothman R, et al. Liability claims and costs before and after implementation of a medical error

- disclosure program. *Ann Intern Med.* 2010;153(4):213–221. PubMed doi: 10.7326/0003-4819-153-4-201008170-00002
73. Adams MA, Elmunzer BJ, Scheiman JM. Effect of a health system's medical error disclosure program on gastroenterology-related claims rates and costs. *Am J Gastroenterol.* 2014;109(4):460–464. PubMed doi: 10.1038/ajg.2013.375
 74. Mello MM, Kachalia A, Roche S, et al. Outcomes in two Massachusetts hospital systems give reason for optimism about communication-and-resolution programs. *Health Aff (Millwood).* 2017;36(10):1795–1803. PubMed doi: 10.1377/hlthaff.2017.0320
 75. Kachalia A, Sands K, Niel MV, et al. Effects of a communication-and-resolution program on hospitals' malpractice claims and costs. *Health Aff (Millwood).* 2018;37(11):1836–1844. PubMed doi: 10.1377/hlthaff.2018.0720
 76. Gallagher TH, Mello MM, Sage WM, Bell SK, McDonald TB, Thomas EJ. Can communication-and-resolution programs achieve their potential? Five key questions. *Health Aff (Millwood).* 2018;37(11):1845–1852. PubMed doi: 10.1377/hlthaff.2018.0727
 77. McMichael BJ, Van Horn RL, Viscusi WK. "Sorry" is never enough: how state apology laws fail to reduce medical malpractice liability risk. *Stanford Law Rev.* 2019;71(2):341–409. PubMed
 78. Berlin L. Will "sorry" make things worse? *Radiology.* 2020;294(1):239–240. PubMed doi: 10.1148/radiol.2019192231
 79. Painter LM, Kidwell KM, Kidwell RP, et al. Do written disclosures of serious events increase risk of malpractice claims? One health care system's experience. *J Patient Saf.* 2018;14(2):87–94. PubMed doi: 10.1097/PTS.0000000000000178
 80. Phillips-Bute B. Transparency and disclosure of medical errors: it's the right thing to do, so why the reluctance? *Campbell Law Rev.* 2013;35(3):332–354.
 81. National Conference of State Legislatures. Medical Professional Apologies Statutes. Accessed July 19, 2024. <https://www.ncsl.org/research/financial-services-and-commerce/medical-professional-apologies-statutes.aspx>
 82. Martin J. An Overview of the Colorado Candor Act. *Colorado Med.* 2019;116(4):53–54.
 83. Dolansky MA, Barg-Walkow L, Barnsteiner J, et al. A call to action following the Radonda Vaught case. *J Nurses Prof Dev.* 2022;38(6):329–332. PubMed doi: 10.1097/NND.0000000000000945
 84. Harrington L. The RaDonda Vaught case: a critical conversation on nursing practice and technology. *AACN Adv Crit Care.* 2023;34(1):11–15. PubMed doi: 10.4037/aacnacc2023873
 85. Institute for Safe Medication Practices. Criminalization of Human Error and a Guilty Verdict: A Travesty of Justice that Threatens Patient Safety. April 7, 2022. Accessed July 19, 2024. <https://psnet.ahrq.gov/issue/criminalization-human-error-and-guilty-verdict-travesty-justice-threatens-patient-safety>
 86. Aubin J, Rivolet O, Taunay AL, Ragot S, Ghazali DA, Oriot D. Benefit of simulation-based training in medical adverse events disclosure in pediatrics. *Pediatr Emerg Care.* 2022;38(2):e622–e627. PubMed doi: 10.1097/PEC.0000000000002454
 87. Stroud L, Wong BM, Hollenberg E, Levinson W. Teaching medical error disclosure to physicians-in-training: a scoping review. *Acad Med.* 2013;88(6):884–892. PubMed doi: 10.1097/ACM.0b013e31828f898f
 88. Amiel JM, Andriole DA, Biskobing DM, et al; Association of American Medical Colleges Core EPAs for Entering Residency Pilot Team. Revisiting the core entrustable professional activities for entering residency. *Acad Med.* 2021;96(7S):S14–S21. PubMed doi: 10.1097/ACM.00000000000004088
 89. AAMC. Core Entrustable Professional Activities for Entering Residency Curriculum Developers' Guide. AAMC; 2014. Accessed April 5, 2024. https://store.aamc.org/downloadable/download/sample/sample_id/63/%20
 90. Lin M, Famiglietti H. Closing the disclosure gap: medical errors in pediatrics. *Pediatrics.* 2019;143(4):e20190221. PubMed doi: 10.1542/peds.2019-0221
 91. Fein S, Hilborne L, Kagawa-Singer M, et al. A conceptual model for disclosure of medical errors. In: Henriksen K, Battles JB, Marks ES, et al, eds. *Advances in Patient Safety: From Research to Implementation (Volume 2: Concepts and Methodology)*. Agency for Healthcare Research and Quality; 2005.

| Situations | Mnemonics and frameworks |
|----------------------------------|---|
| Responding to emotion | <p>N-U-R-S-E-S</p> <ul style="list-style-type: none"> • Naming • Understanding • Respecting • Supporting • Exploring • Silence implementing |
| Building rapport |  |
| Approaching goals of care | <ul style="list-style-type: none"> • Key concepts (know your audience, introduce early, plan to re-discuss) • Restructure the interview (invest in the relationship, intentional body language, voice & tone) • Addressing goals of care (listen, respond to emotion, make a recommendation) |
| Diffusing conflict |  |
| Discussing prognosis | <p>A-D-A-P-T</p> <ul style="list-style-type: none"> • Ask • Discover • Anticipate • Provide • Track emotion |
| Breaking bad news | <p>S-P-I-K-E-S</p> <ul style="list-style-type: none"> • Setup • Perception • Invitation • Knowledge • Empathize • Summarize & strategize |

Collaboratively created by: Wiseman, ML; Wetzel, RA; Fairbrook, SW; Orrick, MJ; Johnson, WR.

In part adapted from VitalTalk© 2019

Disclaimer: The views expressed on this handout are those of the creators and do not necessarily reflect the official policy of the Department of Defense or the U.S. Government.

| Communication Tools | Alternative names & Overlapping tools | Example phrases |
|---------------------------|--|--|
| Anticipate | Normalize, Respect | "Talking about the future can be a little scary." "If you're not sure about <i>[insert topic]</i> , maybe you could tell me how you're thinking about the pros and cons of <i>[insert topic]</i> ." "Uncertainty is difficult and we may know more <i>[insert timeline]</i> ." |
| Ask | Ask-Tell-Ask, Soft Start, Introduce early | "Could we talk about <i>[name topic]</i> ?" "Would it be ok if we talked about <i>[name topic]</i> ?" "If you have <i>[name time]</i> minutes I'd like to ask you about <i>[name topic]</i> ." "What would you like to know about <i>[insert topic, disease, prognosis]</i> ?" "What have other providers told you about your prognosis or future?" |
| Create New Options | | "What would be a tolerable outcome for you?" "Would it be reasonable to trial this..." "Is there someone else that you think we should include to help fix this issue?" "If this plan doesn't work then we can move to the next one." |
| Establish expectations | Plan to revisit/re-discuss, prepare, Introduce early | "This is an ongoing conversation." "Who else would you like to discuss this with." "Let's plan to revisit." "This has the potential to cause <i>[name emotion]</i> , but my goal is to do what is best for <i>[name common goal]</i> " "This is a complex situation and I may appear <i>[name emotion]</i> , could you let me know if that happens?" "I would like to discuss <i>[insert topic]</i> to make sure that we <i>[insert rationale]</i> ." |
| Exploring | Discover | "Tell me more." "Could you say more about what you mean when you say <i>[repeat phrase]</i> " "For some people prognosis is numbers or statistics about how long they will live. For other people, prognosis is about living to a particular date. What would be more helpful for you?" "Some people say that <i>[insert decision, sentiment, opinion, preference/wish]</i> and others say <i>[insert decision, sentiment, opinion, preference/wish]</i> . Which is more like you?" "What have other providers told you about <i>[insert topic]</i> ?" |
| Intentionally Collaborate | Highlight consensus Explore Differences | "I feel like we agree on <i>[name topic/concept/idea]</i> ." "It seems like we have a different view point <i>[name topic/concept/idea]</i> can we explore that more." |
| Invite | Ask, Explore, Understand | "Tell me your thoughts first, then I will describe mine." "I must be missing something, could you help me understand..." "Could you walk me through what you're seeing so that I can better understand your side?" |
| Make a recommendation | Ask, Tell | "Would it be okay if I offered a recommendation?" "I hear that <i>[insert comment]</i> is important to you and <i>[insert comment]</i> would be worse than death. Based on this, I would recommend <i>[insert recommendation]</i> ." |
| Naming | Acknowledge emotion, Notice, Address, Track emotion | "It sounds like you are <i>[name the emotion]</i> ." "It seems like we are interpreting the data differently." "I think we have different points of view and I want to understand yours." "It sounds like this is has had a big impact." "This is difficult." "I wish I had better news." "I can see that this is not what you were hoping for." |

Collaboratively created by: Wiseman, ML; Wetzel, RA; Fairbrook, SW; Orrick, MJ; Johnson, WR.

In part adapted from VitalTalk© 2019

Disclaimer: The views expressed on this handout are those of the creators and do not necessarily reflect the official policy of the Department of Defense or the U.S. Government.

| Communication Tools | Alternative names & Overlapping tools | Example phrases |
|----------------------|---|---|
| Respecting | Normalize, Anticipate | "I can see how hard you've worked to learn about your illness." "You've done a great job taking care of your <i>[loved one]</i> ." "This is a really challenging conversation and we can revisit it. Let's pause for today." "Some people feel <i>[insert emotion]</i> and that is normal." |
| Silence implementing | Body language, Listen | Actively listen with body language (e.g., lean in, nod, sit, keep body open) Comfort with body language (e.g., appropriate touch, maintain eye contact, nod to show listening, lean in) You have two ears and one mouth, use them accordingly (i.e., in that ratio) Avoid interruption |
| Supporting | Empathize, Understanding | "I will do my best to make sure that you have what you need." "This is a tough situation." "I haven't been through that, I can't imagine what it is like." "I'm sorry that this is happening." "Given what has happened I can understand your concern." |
| Tell | Ask-Tell-Ask, Provide, Express transparency | "The worst case scenario is <i>[insert 25th percentile]</i> and the best case scenario is <i>[insert 75th percentile]</i> ." "In 100 people exactly like you, <i>[insert expected outcome]</i> would happen to <i>[insert number]</i> by <i>[insert date]</i> ." From my knowledge of your situation, there is a <i>[insert: good/slim/50-50]</i> chance of <i>[insert outcome with or without timeline]</i> ." "I wish I could say for certain, unfortunately <i>[name limitation]</i> ." |
| Understanding | Supporting | "Given what has happened, I can understand your concern." "You expressing that helps me understand what you are thinking." |
| Voice & tone | | Maintain a calm voice Speak clearly Pause frequently Monitor your speaking cadence (i.e., avoid speaking too fast) |
| Avoid These | | Pithy/Negative personal comments Interjections/unnecessary corrections Yelling/angry tone of voice Disrespectful/closed off body language (arms crossed, eye rolls, standing sideways when talking to someone, shuffling towards the door) |

DO'S AND DONT'S

| Table 3. Mistake disclosure message strategies to avoid | |
|---|---|
| Message strategy | Examples |
| 1. Blocking avenues to questions | "Let's not worry about that now." |
| 2. Redirecting the conversation to less relevant aspects of the mistake | "What I want to focus on is getting better, not what caused the problem." |
| 3. Neglecting to answer questions | "Don't worry about that. Tomorrow we will start treatments." |
| 4. Placing the blame on the patient/family | "Unfortunately, if your weight and diabetes had been under control, it is unlikely this mishap would have happened." |
| 5. Overloading the patient/family with information | "During the operation the bile duct, which carries the bile from the liver down to the gallbladder, was injured because you had inflammation there for so long that I had to peel everything apart, and because of your diabetes you did not heal well, and the bile duct started leaking." |
| 6. Blaming the system | "Because the hospital is under pressure to serve so many patients, we don't have the staffing we need to watch out for these problems. If we had more staff, this mistake would never have happened." |

From "Disclosing Medical Mistakes: A Communication Management Plan for Physicians" (The Permanente Journal/ Spring 2013/ Volume 17 No. 2)

| Table 2. Examples of empathic, exploratory, and validating responses | | |
|--|---|---|
| Empathic statements | Exploratory questions | Validating responses |
| "I can see how upsetting this is to you." | "How do you mean?" | "I can understand how you felt that way." |
| "I can tell you weren't expecting to hear this." | "Tell me more about it." | "I guess anyone might have that same reaction." |
| "I know this is not good news for you." | "Could you explain what you mean?" | "You were perfectly correct to think that way." |
| "I'm sorry to have to tell you this." | "You said it frightened you?" | "Yes, your understanding of the reason for the tests is very good." |
| "This is very difficult for me also." | "Could you tell me what you're worried about?" | "It appears that you've thought things through very well." |
| "I was also hoping for a better result." | "Now, you said you were concerned about your children. Tell me more." | "Many other patients have had a similar experience." |

From "SPIKES--A Six-Step Protocol to Delivering Bad News: Application to the Patient with Cancer" (The Oncologist, 2000)

Sharing Unexpected News

– Set up the interview

Arrange for privacy, involve significant others, sit down, establish rapport with good eye contact, and manage time constraints and interruptions

– Perception of patient/parent

Ask open ended questions to determine what the parent understands about the situation so far, may also allow you to assess medical literacy

– Invitation to information

Obtain agreement from family regarding when, how, and what detail of information they want to know.

– Knowledge delivered

Explain what has happened (e.g. if there was a medical error what the error was) and its impact on the patient.

– Emotional response Summary

Respond to parent's emotional response with empathy. If the bad news involves a medical error, make sure you include an apology.

– and strategy forward

Provide information regarding next steps. In case of ME, what corrective actions are being taken both for the patient and at the systems level to prevent recurrence. If not ME, focus should be on continued care and support for patient and family.

Adapted from "SPIKES-A Six-Step Protocol for Delivering Bad News: Application to the Patient with Cancer" (The Oncologist, 2000)

Disclosure of Medical Error

Establish a rapport and create an appropriate environment for difficult discussion.

Disclose that a medical error occurred.

Apologize for the error.

Answer all parental questions.

Use at least one empathetic statement.

Explain in plain language what has happened and the plan for addressing the situation.

Responding to Emotion (NURSE)

| Step | Example | What you say or do |
|----------------------|--|--|
| Naming | "It sounds like you are frustrated" | In general, turn down the intensity a notch when you name the emotion |
| Understanding | "This helps me understand what you are thinking" | Think of this as another kind of acknowledgment but stop short of suggesting you understand everything (you don't) |
| Respecting | "I can see you have really been trying to follow our instructions" | Remember that praise also fits in here: "I think you have done a great job with this" |
| Supporting | "I will do my best to make sure you have what you need" | Making this kind of commitment is a powerful statement |
| Exploring | "Could you say more about what you mean when you say that..." | Asking a focused question prevents this from seeming too obvious |

Three Fundamental Skills

| Skill | Example | Notes |
|----------------------------|--|--|
| Tell me more | "Tell me more about..." | Use when you are not sure what someone is talking about (rather than jump to an assumption). |
| Ask-tell-ask | "What do you think about..." "Here's what the tests show" "Does that make sense...?" | Related to Assess-Knowledge-Respond in SPIKES. Think of this as one unit of information transfer |
| "I wish" statements | "I wish I could say that the chemo always works" | Enables you to align with the patient while acknowledging the reality of the situation |

Delivering Serious News (GUIDE)

| Step | What you say or do |
|---|---|
| Get Ready – Info, People, Place | <p>“Let me take a minute to make sure I’ve got what I need.”</p> <p>Make sure you have all the information you need at hand.</p> <p>Make sure you have all the right people in the room.</p> <p>Find a place with some privacy.</p> |
| Understand what the patient knows | <p>“What thoughts have you had since the biopsy?”</p> <p>“What have you taken away from other doctors so far?”</p> |
| Inform starting with a headline | <p>“The CT scan shows that the cancer has gotten worse”</p> <p>Give the information clearly and to the point with a one-sentence headline of the most important piece of information you want them to take away.</p> <p>Avoid jargon</p> <p>After the headline you will need to give more information, but after giving the headline, STOP!</p> |
| Demonstrate empathy Respond directly to emotion | <p>“I can see this news is not what you were hoping for.”</p> <p>Expect the patient’s first response to be emotion.</p> <p>Acknowledge the emotion explicitly.</p> |
| Equip the patient for the next step | <p>“Is there anything I could do to make this a little easier?”</p> <p>“I want you to be prepared for the next step. Can I explain...”</p> <p>Don’t dismiss concerns or say that everything will be fine.</p> |

Sharing Unexpected News Quiz


1. According to the Institute of Medicine, patient safety is one of the 6 domains of health care quality, and medical errors are attributable to 44,000 to 98,000 inpatient deaths annually in the United States.

2. What is the difference between an adverse event and a medical error?

Medical Error = commission or omission that unreasonably increases the risk of an undesirable patient outcome.

Adverse event = patient harm due to medical care. Can be preventable (due to medical error) or nonpreventable (not due to medical error.)

3. Complete the labels for the chart:



4. True or False, Physicians have an ethical obligation to disclose AE's and ME's. **TRUE**

5. Which of the following are obstacles to disclosure?

- a. Perceived legal risks
- b. Concern that disclosure will harm patient/family
- c. Fear of embarrassment
- d. Lack of skill in disclosing unexpected/serious news
- e. Language and cultural differences
- f. **All of the above, although legal concerns may be the most significant barrier.**

6. Does Maryland have a protective apology law? DC? Virginia?

YES

[Click Here](#)

7. What are the 6 steps of SPIKES? Spikes was developed for delivering unexpected/serious news in regards to diagnosis and prognosis. Can it be applied to AE's and ME's?

1. **Set up the interview.**

2. **Perception of patient/parent**

3. **Invitation to information.**

4. **Knowledge delivered (medical facts). For cases with MEs, apology should be included**

5. **Emootional response observed and addressed with empathetic statement**

6. **Summary and strategy forward.**

Use this tool as you observe the following scenarios.

Directions: Please indicate whether the physician completed the stated actions, with Y = completed (Yes) or N = did not complete (No)

S – Set the stage

1. Clearly introduced herself/himself
2. Clearly stated his/her role in the care of the patient

P – Perception

3. Determined the level of knowledge the survivors possessed prior to their arrival in the waiting room
4. Took note of the news receiver's vocabulary

I – Inform

5. Briefly indicated the chronology of events leading up to the death of the patient
6. Used language appropriate for the survivor's culture and educational level
7. Avoided using euphemisms

K – Knowledge

8. Allowed the survivor to react to the information and ask questions or express concerns
9. Answered ALL questions in an appropriate manner

E – Empathy

10. Used proper statements to show concern for the grieving
11. Validated emotions of the grieving

S – Summary and Strategy

12. Avoided showing any physician guilt for the loss/poor prognosis
13. Established personal availability to answer questions for the survivor at a later date
14. Ended the discussion and departed in an appropriate manner

"Breaking bad news education for emergency medicine residents: A novel training module using simulation with the SPIKES protocol" From Journal of Emergencies, Trauma and Shock (2010)

Breaking Bad News Case

Please read through the following cases. Take turns in the roles of the provider and the parent. Improvise the discussion around the provided events. If you prefer, you may use one or more of the group's real-life cases to practice disclosing bad news or medical error. Please ensure that you leave time after the improvisation for debrief on what went well, what challenges you faced, and strategies that would improve the disclosure.

1. Spencer Jackson is a 14 year old male who you saw in clinic earlier today due to the complaint of some enlarged lymph nodes in his neck. You detected enlarged fixed nodes in the bilateral cervical chains, and sent Spencer to obtain a chest xray. They have now returned to the clinic after the chest xray and you receive a call from radiology that he has a large mediastinal mass.
2. Dawn Lee is a 3 day old full term AGA female you are seeing in clinic for routine newborn follow-up. Prenatal course was unremarkable, but nursery course was remarkable for DAT+, A neg/O pos incompatibility. Dawn was discharged at 48 hours with a serum bili of 13. Family was offered to stay an extra day for the infant to receive phototherapy, but they declined in favor of close outpatient follow-up. On arrival to the clinic, transcutaneous bili is 16. You discuss the need for a serum bili obtained via heelstick, and the parents are extremely reluctant mentioning that the first heel stick in the hospital caused Dawn terrible pain, but eventually agree. The family lives close by and has returned home. You have had a very busy clinic day and now several hours later, you look up the result and find that the result is "QNS." You are on the phone with Mr. or Ms. Lee to provide them an update.
3. Stephen Tyler is a 17 year old male who was recently discharged from the hospital with a new diagnosis of seizure disorder. During the admission he had a full workup and was started on an antiepileptic medication. He is here now with his parent for hospital follow up and asks you about what he activities should avoid while on the anti-epileptic. You need to counsel him on how he cannot legally drive (in the state of MD) for 3 months from his last seizure.
4. You are seeing John Smith, a 4 year old with asthma for a hospital follow up in continuity clinic 5 days after he was discharged. The patient was recently admitted for an asthma exacerbation and was intended to be sent home with an additional 5 day steroid course, however the ward team did not order it and so the discharge medication was not dispensed. The parents report that he was doing better after going home, but he is now back to requiring albuterol every 4 hours and the patient presents with active wheezing and hypoxia. You will need to disclose the error in discharge medications and provide a plan of care moving forward.
5. You have graduated residency and are made the clinic supervisor at your small MTF. You have a civilian NP who has been at the clinic for years and you are now their supervisor. This individual is continually late in completing their notes, and your leadership is now directing you to counsel them and give them a formal memo for the record as they have 90 notes over the last month that remain unsigned.