



NCC Pediatrics Continuity Clinic Curriculum: **Sharing Unexpected News** *Faculty Version*



Goals & Objectives:

Goal: Increase your knowledge and skill in delivering difficult and unexpected news, and disclosing adverse events to families.

Objectives: At the end of this module, the pediatric resident should be able to:

- Recognize challenges specific to delivering unexpected news and disclosure of medical errors.
- Describe the elements of the mnemonics “SPIKES, NURSE, and GUIDE”
- Apply SPIKES, NURSE, or GUIDE to clinical scenarios that require delivering unexpected news or disclosure of medical errors.

Pre-Meeting Preparation:

- “Disclosure of Adverse Events in Pediatrics” (*AAP Policy Statement, Pediatrics, 2016*)
- "When Bad News Isn't Necessarily Bad: Recognizing Provider Bias When Sharing Unexpected News (*Carroll et al. Pediatrics, 2018*)
- Do's and Dont's, SPIKES strategies for delivering unexpected news
- Watch the [Ted Talk](#) called "Doctors make mistakes. Can we talk about that?"
- Complete quiz questions prior to clinic
- Recall a situation where you had to share unexpected news and be prepared to discuss.

Conference Agenda:

- Review the sharing unexpected news quiz answers (*5 minutes*)
- Discuss your own case when you delivered unexpected news
- Practice delivering unexpected news using the scenarios included here. Make sure that you leave time for meaningful feedback. (*10 minute scenario and 5 minute feedback*)

Extra Credit:

- [Medical Disclosure Powerpoint Slides from Ms. Barbara Moidel](#)
- ["SPIKES -- A Six-Step Protocol for Delivering Bad News: Application to the Patient with Cancer" \(The Oncologist, 2000\)](#)
- [“Disclosing Medical Mistakes: A Communication Management Plan for Physicians" \(Permanente Journal, 2013\)](#)
- [“Teaching Physicians How to Break Bad News” \(Archives of Pediatric and Adolescent Medicine, 1999\)](#)
- ["The Many Faces of Error Disclosure: A Common Set of Elements and a Definition" \(Society of General Internal Medicine, 2007\)](#)
- ["Medical Error Disclosure Among Pediatricians" \(Archives of Pediatric and Adolescent Medicine, 2008\)](#)
- [Institute for Healthcare Improvement Open School Course PS 105: Responding to Adverse Events](#)
- [States with Apology Protection Laws](#)



Disclosure of Adverse Events in Pediatrics

COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT, COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY

Despite increasing attention to issues of patient safety, preventable adverse events (AEs) continue to occur, causing direct and consequential injuries to patients, families, and health care providers. Pediatricians generally agree that there is an ethical obligation to inform patients and families about preventable AEs and medical errors. Nonetheless, barriers, such as fear of liability, interfere with disclosure regarding preventable AEs. Changes to the legal system, improved communications skills, and carefully developed disclosure policies and programs can improve the quality and frequency of appropriate AE disclosure communications.

INTRODUCTION

Patient safety has been characterized as 1 of the 6 domains of health care quality by the Institute of Medicine (IOM),¹ which attributed 44 000 to 98 000 inpatient deaths annually to medical errors (MEs) in the United States.² The American Academy of Pediatrics has called attention to the importance of pediatric patient safety since 2001³ and recommended improved identification and reporting of MEs and adverse events (AEs) to improve the culture of safety in pediatric care.⁴

The IOM defines “patient safety” as the prevention of patient harm and freedom from accidental injury in the health care setting.⁵ An AE occurs when patient harm is caused by medical care.² A ME is an act of commission or omission that unreasonably increases risk of an undesirable patient outcome. AEs may be determined to be preventable (when patient harm is related to a ME) or nonpreventable (when patient harm occurs in the absence of ME). An ME that causes patient harm becomes a preventable AE. An ME that has the potential to cause patient harm but does not do so is referred to as a potential AE or near miss. These concepts and their relationships are illustrated in Fig 1.

The magnitude of harm to patients from AEs can be estimated but has not been quantified reliably. One investigation reported 12.91 AEs per 1000 hospital discharges among patients from birth through 15 years of age,

abstract

FREE

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

DOI: 10.1542/peds.2016-3215

Accepted for publication Sep 26, 2016

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2016 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: The authors have indicated they do not have a financial relationship relevant to this article to disclose.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

To cite: AAP COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT and AAP COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY. Disclosure of Adverse Events in Pediatrics. *Pediatrics*. 2016;138(6):e20163215




FIGURE 1
AEs and MEs.

with “negligence” identified in 28% of events.⁶ In a study of 3700 pediatric hospitalizations, 1% of patients experienced AEs, and 60% of these were determined to be preventable.⁷ Among Canadian inpatient pediatric AEs, the overall rate was 9.2%, and those related to surgery were most frequent.⁸ Almost half of the AEs in this study were determined to be preventable. A prospective study at 6 pediatric practices found that 3% of patients had preventable adverse drug events.⁹ There have been wide variations in estimates of the incidence of AEs in the pediatric inpatient setting and even more uncertainty about the rates of MEs and AEs in the outpatient setting, where most pediatric care occurs.^{4,10} What is clear is that **pediatric AEs are frequent occurrences, with significant patient morbidity and mortality.** The economic cost of AEs was estimated at between \$393 and \$958 billion in 2006, at which time these amounts were equal to 18% to 45% of total US health care spending.¹¹ Specific costs for the subset of cases involving children have not been quantified.

AEs not only affect patients and their families but also may have devastating effects on health care providers, who may suffer emotional consequences both from preventable AEs and from subsequent malpractice litigation.¹² Affected clinicians may feel guilt, shame, and isolation, and these feelings may be exacerbated by negative reactions from their colleagues.^{13–16} Anticipated or actual punitive consequences can add additional emotional and financial burdens on providers.¹⁷

The **concepts of reporting and disclosure of AEs should be distinguished.** “Reporting” refers to the exchange of information among providers and regulators. Reporting systems may be internal to health care organizations or may be required by licensing boards and governmental agencies. These systems may be voluntary or mandatory, and some organizations may use automated AE reporting (eg, bar-code systems and computerized provider order entry). Although in the past, many AE reporting systems focused on punitive consequences, such an approach has been found to deter further reporting. Thus, the current trend has been to reward the reporting of AEs with positive feedback and to use existing reports to develop systematic remedies to promote a safer patient care environment. **“Disclosure,” in contrast, is communication directed from health care personnel to the affected patient and/or family about an AE. Disclosure is a description of what is known and does not include speculation about causation or individuals’ motivations or assumptions about judgment or fault.**

The IOM noted that **most MEs are attributable to flaws in systems rather than individuals** and called for a “dramatic improvement in the reliability and safety” of the health care process.² For this improvement to occur, AEs must first be identified and given attention to understand their preventable causes and to allow for systematic safety improvements. **Disclosure and open communication with patients and their families after an AE may provide benefit to the patient and to the health care provider, reduce consequential harms, allow for better follow-up, and promote a safety culture.** Additionally, patients and caregivers most often desire complete disclosure of AEs and may be less likely to pursue litigation against their health care providers

if complete disclosure is done.¹⁸ Conversely, **lack of communication may make patients feel worse and may erode the sense of trust in their caregivers that is key to healing and to optimal health care.**¹⁸ The health care provider’s self-perception and self-confidence may deteriorate when health care outcomes are poor.^{15,16} Full and open communication may alleviate the provider’s feelings of anxiety, disgrace, and guilt.¹⁹ However, those feelings may also persist and may even be an obstacle to full disclosure. **Pediatric clinicians and institutions should be alert to the involved provider becoming a “second victim” of an ME.**¹⁷

ETHICAL CONSIDERATIONS IN DISCLOSURE

In addition to the quality-control and systems-improvement benefits of disclosure, physicians broadly acknowledge that **disclosure of AEs is an ethical obligation.**^{20–23} Although there is evidence that they often do not “practice what they preach,”^{13,18,24–27} physicians and medical trainees, in particular, agree that physicians have an ethical obligation to their patients to disclose preventable AEs.^{21,26–28} In an anonymous survey among pediatric residents and attending pediatricians, pediatricians in private practice were less likely to report errors than other attending pediatricians (72% vs 92%; $P < .001$).²⁵ Most agreed that disclosing a serious error would be difficult (overall, 88%; attending pediatricians, 86%; residents, 96%; $P = .005$). More residents than attending pediatricians had received education about how to disclose errors (57% vs 29%; $P < .001$).

BARRIERS TO DISCLOSURE

Despite the compelling benefits and ethical imperatives for AE disclosure, many physicians nevertheless continue to have difficulty completing the task of informing

patients and their families about AEs and MEs.^{13,21,25,29} Several barriers can create obstacles to disclosure, including perceived legal risks and the cautionary advice of legal counsel, concerns that disclosure might harm patients, a lack of confidence in disclosure skills, and a fear of embarrassment.^{14,25,30-32} Language and cultural differences may also interfere with meaningful communication about AEs.¹⁴ Among these barriers, perceived legal risks and legal advice cautioning against disclosure may be the most significant.^{27,32}

LEGAL RISKS ASSOCIATED WITH DISCLOSURE AND APOLOGY

Historically, lawyers advised their physician-clients not to disclose MEs and did so with sound legal justification.³³⁻³⁶ In the US legal system, previous statements by an observer of an event generally are not permitted to be introduced in court as evidence about that event.³⁷ In other words, a courtroom witness can testify about what he knows from his own observation but not about what someone else told him about the event. This “hearsay rule” presumes that bringing a witness into the courtroom to discuss what he actually knows from firsthand experience is more informative and more reliable than having someone else recount what another observer, on some previous occasion, said that she had seen.³⁷ However, the law recognizes the possibility that a witness might be less likely to fully and truthfully describe his observations when that witness has become the defendant in the lawsuit. Moreover, that defendant-witness might recant truthful statements that he might have previously made about the event. Therefore, a well-established exception to the hearsay rule permits previous, self-incriminating statements by defendants to be admitted into evidence, even though those

statements are hearsay.³⁸ This “admission of a party-opponent” exception has been used by plaintiffs in numerous cases to quote a defendant-physician’s earlier admission of fault to help prove that the physician-defendant committed malpractice.³⁶

The concept of “apology” generates similar legal concerns but may be even more problematic because of variations in definition. “Apology” is defined by *Webster’s Dictionary* as “a statement that you are sorry about something” or “an expression of regret for having done or said something wrong.”³⁹ This definition does not include any acknowledgment of fault. Alternatively, “apology” may be defined as “an admission of error or discourtesy accompanied by an expression of regret.” This lack of clarity in what it means “to apologize” can result in plaintiffs asserting in court that a physician-defendant “admitted” to fault, when the physician-defendant had no intention of admitting fault in the course of the apology.⁴⁰

Some lawmakers have attempted to encourage physician disclosure and apology by reducing this legal risk through legislative changes to the hearsay rule exceptions.^{30,33,41} Many state legislatures have attempted to encourage disclosures and apologies by adopting “apology laws,” intended to encourage physician-patient communications about AEs by limiting, or even completely excluding, plaintiffs’ ability to use such communications as evidence against the physician in later litigation.^{30,41} “Sympathy-only” apology laws bar the use of physician expressions of sympathy by plaintiffs at trial to prove negligence but do permit plaintiffs to use physician statements that expressly admitted fault.³⁰ On the other hand, “admission of fault” apology laws prevent plaintiffs from using virtually any previous physician disclosures to the

patient and family, even when the physician admitted responsibility for an ME or other improper care, to prove negligence at trial.^{30,*}

The effects of these apology laws in reducing liability risks remain unclear. Nevertheless, there is some evidence that they may be effective in reducing liability risks. For example, in *Deutsch v INOVA Health Care Services*, the plaintiff alleged that the defendant cardiologist was negligent in failing to personally evaluate a tachycardic patient in the emergency department.⁴² The cardiologist subsequently met with the family in the ICU and reportedly stated, “I am sorry I wasn’t there.”⁴² The Circuit Court of Virginia prohibited the plaintiffs from using this statement at trial, declaring that this “expression of sympathy and benevolence” was precisely the type of communication that the Virginia apology law was enacted to protect.⁴² Similarly, in *Airasian v Shaak*, the Georgia Court of Appeals prohibited the plaintiffs from using as evidence the defendant surgeon’s postoperative statement to the patient’s wife that “this [need for a second, emergency surgery] was my fault.”⁴³

The *Deutsch* and *Airasian* cases demonstrate how apology laws can protect physician disclosures about AEs, thus encouraging other physicians to communicate with patients and potentially preserving physician-patient relationships, improving patient and family satisfaction, and helping to identify systems weaknesses leading to AEs. However, some apology laws, particularly sympathy-only apology laws, may not provide protection and reassurance sufficient to encourage disclosure.^{33,36} Mastroianni et al opined that “sympathy-only” apology laws have a fatal structural weakness in that they do not protect the key

*For information on current state apology laws, please contact the AAP Division of State Government Affairs at 800/433-9016, extension 7799, or at stgov@aap.org.

information that patients want communicated to them after an AE.³³ These authors observed that **patients (and families) want full and clear explanations and accountability for adverse outcomes. Mere expressions of sympathy, in the absence of full disclosure and accountability, may be unsatisfactory to patients and may fail to bring about systems improvements that might prevent future adverse outcomes.**¹³

There is evidence that disclosure, even in the absence of protective apology laws, may reduce liability risks and litigation costs.^{44,45} Nevertheless, some researchers have suggested a theoretical risk that the sheer volume of patients who could be put on notice of MEs by full disclosure policies might result in a greater volume of lawsuits.⁴⁶

FACILITATING BETTER DISCLOSURE

In the event of a preventable AE, the **pediatrician's first responsibility is to attend to the immediate medical needs of the patient.** Thereafter, **improved physician-patient communication may contribute to enhanced patient satisfaction, better ongoing medical care for the patient, and prevention of future AEs.** Moreover, full disclosure about such events is **ethically indicated.** Therefore, pediatricians should endeavor to provide appropriate disclosure when their patients experience preventable AEs. Pediatricians can help facilitate such disclosures in at least 2 ways: developing and implementing disclosure policies and procedures for their own practices, and supporting public policies that facilitate disclosure.

Unfortunately, preventable AEs are often the result of a cascade of errors rather than a single mistake.⁴⁷ **Gaining a clear understanding of whether an ME occurred, how it occurred, and whether and how it affected a patient may be a**

difficult and complex process. Hasty confessions have the potential to generate legally admissible admissions of fault, even if the confession later turns out to have been erroneous. To reduce the risks of inaccurate or premature admissions of fault, practices and hospitals may choose to establish disclosure policies that govern how preventable AEs are investigated and how the findings of such investigations are communicated to patients and families.

Pediatricians can prepare for future AEs by developing disclosure policies and procedures. Disclosure plans will help pediatricians determine in advance what types of information that they will communicate with patients and families when AEs occur. Pediatricians can be ready to appropriately express their sympathy about adverse outcomes and to declare their willingness and intention to investigate the cause(s) of the AE, to take measures (when appropriate) to prevent similar events in the future, and to provide ongoing necessary medical care and support to the affected patient and family. Pediatricians can also be prepared to take appropriate steps to protect themselves from unnecessary and inappropriate medical liability, by having a plan in place to notify medical malpractice insurance carriers and to consult with legal counsel.

EDUCATION ON DISCLOSURE OF PREVENTABLE AEs

Education regarding practical disclosure skills and patient safety can be valuable to pediatric trainees and clinicians at all levels of experience. Simulation technology has been successfully used as one technique to assist in this learning.⁴⁸ Several other approaches to resident patient safety curricula have been reported.⁴⁹

CONCLUSIONS

There is little doubt that patients, families, and physicians are better served when full and honest communications can take place after AEs. The American Academy of Pediatrics and its members can help to promote such communications by becoming informed about, and encouraging, state apology laws and other public policies that support disclosure.

RECOMMENDATIONS

1. Pediatric health care providers and institutions should develop and implement their own policies and procedures for identifying and disclosing AEs to patients and families in an honest and empathetic manner as part of a nonpunitive culture of ME reporting.
2. Pediatric institutions and practices should develop policies and procedures to provide emotional support for clinicians involved in AEs.
3. Pediatric medical educators should develop and implement educational programs regarding identification and prevention of MEs and communication about AEs with patients and their families as part of a comprehensive patient safety curriculum.
4. Additional research is needed on the consequences of various approaches to disclosure as well as of the effectiveness of disclosure education.
5. State legislators and other governmental and regulatory bodies are encouraged to continue developing apology laws and other mechanisms to reduce liability risks associated with disclosure.

LEAD AUTHORS

William M. McDonnell, MD, JD, FAAP
Daniel R. Neuspiel, MD, MPH, FAAP

COMMITTEE ON MEDICAL LIABILITY AND RISK MANAGEMENT, 2015–2016

William M. McDonnell, MD, JD, FAAP, Chairperson
Robin L. Altman, MD, FAAP
Steven A. Bondi, JD, MD, FAAP
Jon Mark Fanaroff, MD, JD, FAAP
Sandeep K. Narang, MD, JD, FAAP
Richard L. Oken, MD, FAAP
John W. Rusher, MD, JD, FAAP
Karen A. Santucci, MD, FAAP
James P. Scibilia, MD, FAAP
Susan M. Scott, MD, JD, FAAP

STAFF

Julie Kersten Ake

COUNCIL ON QUALITY IMPROVEMENT AND PATIENT SAFETY, 2014–2015

Wayne H. Franklin, MD, MPH, MMM, FAAP, Chairperson
Terry Adirim, MD, MPH, FAAP
David Gordon Bundy, MD, MPH, FAAP
Laura Elizabeth Ferguson, MD, FAAP
Sean Patrick Gleeson, MD, MBA, FAAP
Michael Leu, MD, MS, MHS, FAAP
Brigitta U. Mueller, MD, MHCM, FAAP
Daniel Robert Neuspiel, MD, MPH, FAAP
Michael Lawrence Rinke, MD, PhD, FAAP
Richard N. Shiffman, MD, MCIS, FAAP
Joel Sanford Tieder, MD, MPH, FAAP

STAFF

Lisa Krams, MS

ABBREVIATIONS

AE: adverse event
IOM: Institute of Medicine
ME: medical error

REFERENCES

1. Institute of Medicine, Committee on Quality of Health Care in America. *Crossing the Quality Chasm: A New Health System for the 21st Century*. Washington, DC: National Academy Press; 2001
2. Kohn LT, Corrigan JM, Donaldson MS, eds. *To Err Is Human: Building a Safer Health System*. Washington, DC: National Academy Press; 2000
3. Lannon CM, Coven BJ, Lane France F, et al; National Initiative for Children's Health Care Quality Project Advisory Committee. Principles of patient safety in pediatrics. *Pediatrics*. 2001;107(6):1473–1475
4. Miller MR, Takata G, Stucky ER, Neuspiel DR; Steering Committee on Quality Improvement and Management and Committee on Hospital Care. Policy statement—principles of pediatric patient safety: reducing harm due to medical care. *Pediatrics*. 2011;127(6):1199–1210
5. Aspden P, Corrigan J, Wolcott J, et al, eds; Institute of Medicine, Committee on Data Standards for Patient Safety. *Patient Safety: Achieving a New Standard for Care*. Washington, DC: National Academies Press; 2004
6. Leape LL, Brennan TA, Laird N, et al. The nature of adverse events in hospitalized patients. Results of the Harvard Medical Practice Study II. *N Engl J Med*. 1991;324(6):377–384
7. Woods D, Thomas E, Holl J, Altman S, Brennan T. Adverse events and preventable adverse events in children. *Pediatrics*. 2005;115(1):155–160
8. Matlow AG, Baker GR, Flintoft V, et al. Adverse events among children in Canadian hospitals: the Canadian Paediatric Adverse Events Study. *CMAJ*. 2012;184(13):E709–E718
9. Kaushal R, Goldmann DA, Keohane CA, et al. Adverse drug events in pediatric outpatients. *Ambul Pediatr*. 2007;7(5):383–389
10. Neuspiel DR, Stubbs EH. Patient safety in ambulatory care. *Pediatr Clin North Am*. 2012;59(6):1341–1354
11. Goodman JC, Villarreal P, Jones B. The social cost of adverse medical events, and what we can do about it. *Health Aff (Millwood)*. 2011;30(4):590–595
12. Wu AW. Medical error: the second victim. The doctor who makes the mistake needs help too. *BMJ*. 2000;320(7237):726–727
13. Gallagher TH, Waterman AD, Ebers AG, Fraser VJ, Levinson W. Patients' and physicians' attitudes regarding the disclosure of medical errors. *JAMA*. 2003;289(8):1001–1007
14. Berlinger N, Wu AW. Subtracting insult from injury: addressing cultural expectations in the disclosure of medical error. *J Med Ethics*. 2005;31(2):106–108
15. Abd Elwahab S, Doherty E. What about doctors? The impact of medical errors. *Surgeon*. 2014;12(6):297–300
16. Waterman AD, Garbutt J, Hazel E, et al. The emotional impact of medical errors on practicing physicians in the United States and Canada. *Jt Comm J Qual Patient Saf*. 2007;33(8):467–476
17. Marmon LM, Heiss K. Improving surgeon wellness: the second victim syndrome and quality of care. *Semin Pediatr Surg*. 2015;24(6):315–318
18. Helmchen LA, Richards MR, McDonald TB. How does routine disclosure of medical error affect patients' propensity to sue and their assessment of provider quality? Evidence from survey data. *Med Care*. 2010;48(11):955–961
19. Bell SK, Mann KJ, Truog R, Lantos JD. Should we tell parents when we've made an error? *Pediatrics*. 2015;135(1):159–163
20. Bernstein M, Brown B. Doctors' duty to disclose error: a deontological or Kantian ethical analysis. *Can J Neuro Sci*. 2004;31(2):169–174
21. Gallagher TH, Waterman AD, Garbutt JM, et al. US and Canadian physicians' attitudes and experiences regarding disclosing errors to patients. *Arch Intern Med*. 2006;166(15):1605–1611
22. Stokes SL, Wu AW, Pronovost PJ. Ethical and practical aspects of disclosing adverse events in the emergency department. *Emerg Med Clin North Am*. 2006;24(3):703–714
23. Waite M. To tell the truth: the ethical and legal implications of disclosure of medical error. *Health Law J*. 2005;13:1–33
24. Blendon RJ, DesRoches CM, Brodie M, et al. Views of practicing physicians and the public on medical errors. *N Engl J Med*. 2002;347(24):1933–1940
25. Garbutt J, Brownstein DR, Klein EJ, et al. Reporting and disclosing medical errors: pediatricians' attitudes and behaviors. *Arch Pediatr Adolesc Med*. 2007;161(2):179–185
26. Cole AP, Block L, Wu AW. On higher ground: ethical reasoning and its relationship with error disclosure. *BMJ Qual Saf*. 2013;22(7):580–585
27. Varjavand N, Bachegowda LS, Gracely E, Novack DH. Changes in intern attitudes toward medical error and disclosure. *Med Educ*. 2012;46(7):668–677

28. Parker M. A fair dinkum duty of open disclosure following medical error. *J Law Med.* 2012;20(1):35–43
29. Gallagher TH, Studdert D, Levinson W. Disclosing harmful medical errors to patients. *N Engl J Med.* 2007;356(26):2713–2719
30. McDonnell WM, Guenther E. Narrative review: do state laws make it easier to say “I’m sorry?” *Ann Intern Med.* 2008;149(11):811–816
31. Gallagher THA. A 62-year-old woman with skin cancer who experienced wrong-site surgery: review of medical error. *JAMA.* 2009;302(6):669–677
32. Lamb RM, Studdert DM, Bohmer RMJ, Berwick DM, Brennan TA. Hospital disclosure practices: results of a national survey. *Health Aff (Millwood).* 2003;22(2):73–83
33. Mastroianni AC, Mello MM, Sommer S, Hardy M, Gallagher TH. The flaws in state “apology” and “disclosure” laws dilute their intended impact on malpractice suits. *Health Aff (Millwood).* 2010;29(9):1611–1619
34. Hyman DA. When and why lawyers are the problem. *DePaul L Rev.* 2008;57(2):267–280
35. Tabler N. Should physicians apologize for medical errors? *Health Lawyer.* 2007;19:23–30
36. Raper SE. No role for apology: remedial work and the problem of medical injury. *Yale J Health Policy Law Ethics.* 2011;11(2):267–330
37. Broun KS, Dix GE, Imwinkelried DJ, et al. The hearsay rule and its exceptions: the hearsay rule. In: Broun KS, Dix GE, Imwinkelried DJ, et al, eds. *McCormick on Evidence.* 7th ed. Eagan, MN: Thomson Reuters; 2013:175–257
38. Broun KS, Dix GE, Imwinkelried DJ, et al. The hearsay rule and its exceptions: admissions of a party-opponent. In: Broun KS, Dix GE, Imwinkelried DJ, et al, eds. *McCormick on Evidence.* 7th ed. Eagan, MN: Thomson Reuters; 2013:259–350
39. *Merriam-Webster Dictionary.* Available at: www.merriam-webster.com/dictionary/apology. Accessed May 24, 2016
40. *Johnson v Randall Smith*, 989 NE2d 35 (Ohio 2013)
41. Clinton HR, Obama B. Making patient safety the centerpiece of medical liability reform. *N Engl J Med.* 2006;354(21):2205–2208
42. *Deitsch v INOVA Health Care Services*, 2005 WL 4876742, No. 223119 (Va Cir Ct 2005)
43. *Airasian v Shaak*, 289 Ga App 540, 657 S.E.2d 600 (2008)
44. Kachalia A, Kaufman SR, Boothman R, et al. Liability claims and costs before and after implementation of a medical error disclosure program. *Ann Intern Med.* 2010;153(4):213–221
45. Wu AW, Huang IC, Stokes S, Pronovost PJ. Disclosing medical errors to patients: it’s not what you say, it’s what they hear. *J Gen Intern Med.* 2009;24(9):1012–1017
46. Studdert DM, Mello MM, Gawande AA, Brennan TA, Wang YC. Disclosure of medical injury to patients: an improbable risk management strategy. *Health Aff (Millwood).* 2007;26(1):215–226
47. Reason J. Human error: models and management. *BMJ.* 2000;320(7237):768–770
48. Matos FM, Raemer DB. Mixed-realism simulation of adverse event disclosure: an educational methodology and assessment instrument. *Simul Healthc.* 2013;8(2):84–90
49. Neuspiel DR, Hyman D, Lane M. Quality improvement and patient safety in the pediatric ambulatory setting: current knowledge and implications for residency training. *Pediatr Clin North Am.* 2009;56(4):935–951

DO'S AND DONT'S

Table 3. Mistake disclosure message strategies to avoid	
Message strategy	Examples
1. Blocking avenues to questions	"Let's not worry about that now."
2. Redirecting the conversation to less relevant aspects of the mistake	"What I want to focus on is getting better, not what caused the problem."
3. Neglecting to answer questions	"Don't worry about that. Tomorrow we will start treatments."
4. Placing the blame on the patient/family	"Unfortunately, if your weight and diabetes had been under control, it is unlikely this mishap would have happened."
5. Overloading the patient/family with information	"During the operation the bile duct, which carries the bile from the liver down to the gallbladder, was injured because you had inflammation there for so long that I had to peel everything apart, and because of your diabetes you did not heal well, and the bile duct started leaking."
6. Blaming the system	"Because the hospital is under pressure to serve so many patients, we don't have the staffing we need to watch out for these problems. If we had more staff, this mistake would never have happened."

From "Disclosing Medical Mistakes: A Communication Management Plan for Physicians" (The Permanente Journal/ Spring 2013/ Volume 17 No. 2)

Table 2. Examples of empathic, exploratory, and validating responses		
Empathic statements	Exploratory questions	Validating responses
"I can see how upsetting this is to you."	"How do you mean?"	"I can understand how you felt that way."
"I can tell you weren't expecting to hear this."	"Tell me more about it."	"I guess anyone might have that same reaction."
"I know this is not good news for you."	"Could you explain what you mean?"	"You were perfectly correct to think that way."
"I'm sorry to have to tell you this."	"You said it frightened you?"	"Yes, your understanding of the reason for the tests is very good."
"This is very difficult for me also."	"Could you tell me what you're worried about?"	"It appears that you've thought things through very well."
"I was also hoping for a better result."	"Now, you said you were concerned about your children. Tell me more."	"Many other patients have had a similar experience."

From "SPIKES--A Six-Step Protocol to Delivering Bad News: Application to the Patient with Cancer" (The Oncologist, 2000)

Sharing Unexpected News

— Set up the interview

Arrange for privacy, involve significant others, sit down, establish rapport with good eye contact, and manage time constraints and interruptions

— Perception of patient/parent

Ask open ended questions to determine what the parent understands about the situation so far, may also allow you to assess medical literacy

— Invitation to information

Obtain agreement from family regarding when, how, and what detail of information they want to know.

— Knowledge delivered

Explain what has happened (e.g. if there was a medical error what the error was) and its impact on the patient.

— Emotional response Summary

Respond to parent's emotional response with empathy. If the bad news involves a medical error, make sure you include an apology.

— and strategy forward

Provide information regarding next steps. In case of ME, what corrective actions are being taken both for the patient and at the systems level to prevent recurrence. If not ME, focus should be on continued care and support for patient and family.

Adapted from "SPIKES-A Six-Step Protocol for Delivering Bad News: Application to the Patient with Cancer" (The Oncologist, 2000)

Disclosure of Medical Error

Establish a rapport and create an appropriate environment for difficult discussion.

Disclose that a medical error occurred.

Apologize for the error.

Answer all parental questions.

Use at least one empathetic statement.

Explain in plain language what has happened and the plan for addressing the situation.

Responding to Emotion (NURSE)

Step	Example	What you say or do
Naming	"It sounds like you are frustrated"	In general, turn down the intensity a notch when you name the emotion
Understanding	"This helps me understand what you are thinking"	Think of this as another kind of acknowledgment but stop short of suggesting you understand everything (you don't)
Respecting	"I can see you have really been trying to follow our instructions"	Remember that praise also fits in here: "I think you have done a great job with this"
Supporting	"I will do my best to make sure you have what you need"	Making this kind of commitment is a powerful statement
Exploring	"Could you say more about what you mean when you say that..."	Asking a focused question prevents this from seeming too obvious

Three Fundamental Skills

Skill	Example	Notes
Tell me more	"Tell me more about..."	Use when you are not sure what someone is talking about (rather than jump to an assumption).
Ask-tell-ask	"What do you think about..." "Here's what the tests show" "Does that make sense...?"	Related to Assess-Knowledge-Respond in SPIKES. Think of this as one unit of information transfer
"I wish" statements	"I wish I could say that the chemo always works"	Enables you to align with the patient while acknowledging the reality of the situation

Delivering Serious News (GUIDE)

Step	What you say or do
<p>Get Ready – Info, People, Place</p>	<p>“Let me take a minute to make sure I’ve got what I need.”</p> <p>Make sure you have all the information you need at hand.</p> <p>Make sure you have all the right people in the room.</p> <p>Find a place with some privacy.</p>
<p>Understand what the patient knows</p>	<p>“What thoughts have you had since the biopsy?”</p> <p>“What have you taken away from other doctors so far?”</p>
<p>Inform starting with a headline</p>	<p>“The CT scan shows that the cancer has gotten worse”</p> <p>Give the information clearly and to the point with a one-sentence headline of the most important piece of information you want them to take away.</p> <p>Avoid jargon</p> <p>After the headline you will need to give more information, but after giving the headline, STOP!</p>
<p>Demonstrate empathy Respond directly to emotion</p>	<p>“I can see this news is not what you were hoping for.”</p> <p>Expect the patient’s first response to be emotion.</p> <p>Acknowledge the emotion explicitly.</p>
<p>Equip the patient for the next step</p>	<p>“Is there anything I could do to make this a little easier?”</p> <p>“I want you to be prepared for the next step. Can I explain...”</p> <p>Don’t dismiss concerns or say that everything will be fine.</p>

Sharing Unexpected News Quiz


1. According to the Institute of Medicine, **patient safety** is one of the 6 domains of health care quality, and **medical errors** are attributable to 44,000 to 98,000 inpatient deaths annually in the United States.

2. What is the difference between an adverse event and a medical error?

Medical Error = commission or omission that unreasonably increases the risk of an undesirable patient outcome.

Adverse event = patient harm due to medical care. Can be preventable (due to medical error) or nonpreventable (not due to medical error.)

3. Complete the labels for the chart:



4. True or False, Physicians have an ethical obligation to disclose AE's and ME's. **TRUE**

5. Which of the following are obstacles to disclosure?

- Perceived legal risks
- Concern that disclosure will harm patient/family
- Fear of embarrassment
- Lack of skill in disclosing unexpected/serious news
- Language and cultural differences
- All of the above, although legal concerns may be the most significant barrier.**

6. Does Maryland have a protective apology law? DC? Virginia?

YES

[Click Here](#)

7. What are the 6 steps of SPIKES? Spikes was developed for delivering unexpected/serious news in regards to diagnosis and prognosis. Can it be applied to AE's and ME's?

- Set up the interview.**
- Perception of patient/parent**
- Invitation to information.**
- Knowledge delivered (medical facts). For cases with MEs, apology should be included**
- Emotional response observed and addressed with empathetic statement**
- Summary and strategy forward.**

Use this tool as you observe the following scenarios.

Directions: Please indicate whether the physician completed the stated actions, with Y = completed (Yes) or N = did not complete (No)

S – Set the stage

1. Clearly introduced herself/himself
2. Clearly stated his/her role in the care of the patient

P – Perception

3. Determined the level of knowledge the survivors possessed prior to their arrival in the waiting room
4. Took note of the news receiver's vocabulary

I – Inform

5. Briefly indicated the chronology of events leading up to the death of the patient
6. Used language appropriate for the survivor's culture and educational level
7. Avoided using euphemisms

K – Knowledge

8. Allowed the survivor to react to the information and ask questions or express concerns
9. Answered ALL questions in an appropriate manner

E – Empathy

10. Used proper statements to show concern for the grieving
11. Validated emotions of the grieving

S – Summary and Strategy

12. Avoided showing any physician guilt for the loss/poor prognosis
13. Established personal availability to answer questions for the survivor at a later date
14. Ended the discussion and departed in an appropriate manner

“Breaking bad news education for emergency medicine residents: A novel training module using simulation with the SPIKES protocol” From Journal of Emergencies, Trauma and Shock (2010)

Sharing Unexpected News Case

1. Please share any personal cases you have experienced in disclosing medical errors/ adverse events as well as in delivering unexpected news. What was difficult? What went well? How would you do things differently next time?

2. Please read through the following cases. Take turns in the roles of the provider and the parent. Improvise the discussion around the provided events. If you prefer, you may use one or more of the group's real-life cases to practice disclosing serious news or medical error instead. Please ensure that you leave time after the parent/provider discussion for a debrief on what went well, what challenges you faced, and strategies that would improve the disclosure.

1. Forsythia Lee is a 3 day old full term AGA female you are seeing in clinic for routine newborn follow-up. Her prenatal course was unremarkable, but nursery course was remarkable for DAT+, A neg/O pos incompatibility. Forsythia was discharged at 48 hours of life with a serum bili of 13. The nursery team recommended Forsythia stay an extra day to receive phototherapy, but her parents declined in favor of close outpatient follow-up. On arrival to the clinic, Forsythia is visibly jaundiced to at least her abdomen. Transcutaneous bili is 16. You discuss the need for a serum bilirubin obtained via heelstick. The parents are extremely reluctant, mentioning that the first heel stick in the hospital caused Forsythia terrible pain. They eventually agree. The family lives close by and has returned home. You have had a very busy clinic day and now as the clinic is closing, you look up the result and find that the lab was canceled due to "quantity insufficient." You are on the phone with Mr. or Ms. Lee to provide them an update. *Identify a resident each to play Mr/s. Lee and the physician. Let the scenario run no longer than 10 minutes, allowing at least 5 minutes for feedback (15 minutes total.) Have the other attendees take notes regarding adherence to SPIKES format, inclusion of apology for error (reported insufficient sample and delay in checking lab.) Ensure a supportive environment for the resident playing physician and have fun.*

2. Forsythia Lee is now 7 days old. She eventually had her bilirubin repeated, although her family failed to understand the urgency of the situation, was overwhelmed with other obligations, and returned for repeat labs at 90 hours of life. On repeat presentation, Forsythia's serum bili was 30. She was admitted to the NICU and is s/p exchange transfusion. She has been noted to have poor feeding and hypotonia and over the past 24 hours has developed extensor muscle hypertonia with opisthotonos. Forsythia's parents note that she seems different today. They are unsure of how to interpret the change they notice and have asked you for an update. You have just finished rounds during which it was discussed that Forsythia most likely has bilirubin encephalopathy which you know to be associated with poor long term developmental outcomes. You are feeling guilty about the initial delayed lab result and your role in this outcome and have continued to have challenging interactions with the Lee family.

As above, identify a resident to play the physician and a resident to play the parent. This scenario involves both medical error and a significant poor outcome. Leave time for feedback!